

Study of Thyroid Function in Patients with Chronic Kidney Disease

Dr. J. Puneekar¹, Dr. A. A. Singh², Dr. Manoj Kumar Malav³

¹Associate Professor, ²Assistant Professor, ³Post Graduate Student,
Department of General Medicine, NSCB Medical College, Jabalpur.

Corresponding Author: Dr. Janki Puneekar

Received: 16/12/2016

Revised: 06/01/2017

Accepted: 09/01/2017

ABSTRACT

Background: Thyroid dysfunction is a commonly seen endocrine abnormality among CKD patients. CKD has been known to affect the pituitary-thyroid axis and peripheral metabolism of thyroid hormones. This study was conducted to find out possible association of CKD and thyroid dysfunction and to estimate the occurrence of thyroid dysfunction in patients with chronic kidney disease and its correlation with severity of renal disease.

Materials and methods: A case control study was designed to study association of CKD cases and thyroid dysfunction. Total 75 CKD cases were recruited in case group and these cases were investigated, serum thyroid level measured by radioimmunoassay and their results were compared to a control group of 75 age/sex matched healthy subject.

Observations: Results of this study showed that majority of the cases were 30-59 years age group (48.21 ±13.7). The male to female ratio in case group was 1.08:1 and in control group 1.02:1. Thyroid dysfunction was found to have 80% more occurrence in CKD Patients as compared to normal healthy controls. Most common thyroid dysfunction observed was overt hypothyroidism (42.66%) followed by Low T₃ syndrome (21.33%), subclinical hypothyroidism (13.33%) and hyperthyroidism (2.7%). With the increasing severity of CKD, overt hypothyroidism also increased i.e. from 16.66% in stage-III, 38.8% in stage-IV to 47.06% in Stage-V. There was significant reduction of serum T₃ level (69.58 ± 38.28ng/dl), T₄ level (6.07± 2.55µg/dl) and elevation of TSH level (7.42± 4.25µIU/ml) in cases as compared to controls (p<0.001).

Conclusion: Thyroid dysfunction (overt hypothyroidism) showed a significant and strong correlation with CKD cases and various stage of CKD. Thus timely identification and treatment of these thyroid abnormalities could alter the course of the disease and help in reducing morbidity of CKD patients.

Keywords: thyroid function, chronic kidney disease.

INTRODUCTION

Thyroid hormones are necessary for growth and development of the kidney and for maintenance of water and electrolyte homeostasis. On the other hand, Kidney plays an important role in the metabolism, degradation and excretion of thyroid hormone. The kidney normally contributes to the clearance of iodine, primarily by glomerular filtration. Thus iodide excretion

is diminished in advanced renal failure, leading sequentially to an elevated plasma inorganic iodide concentration and an initial increment in thyroidal iodide uptake. [1] Increased total body inorganic iodide can potentially block thyroid hormone production by affecting the pituitary-thyroid axis and peripheral metabolism of thyroid hormones. Such changes explain higher frequency of hypothyroidism in patients

with chronic kidney disease. So, Thyroid dysfunction is a commonly seen endocrine abnormality among CKD patients. [2]

It has been shown that in chronic kidney disease (CKD), as the glomerular filtration rate (GFR) falls, there is a higher possibility of developing clinical and subclinical hypothyroidism (SCH). [3] Prevalence of Thyroid dysfunction in CKD is found to be ranging from 13% in early CKD to 70% in ESRD according to various studies. [4-7] The relation between thyroid dysfunction and severity of CKD is not clear. Several previous studies depict conflicting results both positive and negative. Thus, there are huge numbers of patients remaining to be diagnosed and/or treated. The present study was conducted to find out possible association of CKD and thyroid dysfunction and to estimate the occurrence of thyroid dysfunction in patients with chronic kidney disease and its correlation with severity of renal disease and also see the clinical presentation of chronic kidney disease patients with thyroid dysfunction or without thyroid dysfunction.

MATERIALS AND METHODS

It is Hospital based Cross sectional study conducted in the department of Medicine at N.S.C.B MCH, Jabalpur (M.P.). A total of 150 patients included in this study out of which 75 patients were chronic kidney disease and 75 were normal healthy patients. The study period extended from March 2015 to August 2016.

Patients with associated history of acute illness, thyroidectomy/thyroid disorder recent surgery, trauma or burns, drugs altering Thyroid profile (like amiodarone, steroids, phenytoin, beta-blockers, estrogen pills, iodine containing drugs, children's and ANC Patients were excluded from the study.

All the subjects i.e. both cases & controls were subjected to medical examination as per a fixed proforma. Morning sample blood was drawn after 12 hrs fasting. The samples of blood were allowed to stand to clot. Serum was

separated by centrifugation, and analyzed by the following methods. Serum Urea Estimated by Diacetyl Monoxide Method (DAM, Method), serum creatinine is estimated by Jaffe's method, and Estimation of T₃, T₄ & TSH by Chemiluminescent immunoassay (CLIA) method. Thyroid dysfunction was considered if patient's thyroid hormones level fall outside the reference range. Expected range of TT₃ is 70-204 ng/dl, expected range of TT₄ is 4.5-11microgm/dl and expected ranges of values of TSH is 0.5-4.5mU/L.

Data analysis

The data was fed into an excel spreadsheet and then tabulated. Data was statistically analyzed using t-test, chi-square test, Fisher's exact test using SPSS and Microsoft excel. p<0.05 was considered to be statistically significant.

RESULTS

The study population comprised 75 CKD patients and 75 normal healthy controls with 39 (52%) males and 36 (48%) females in case group and 38(50.7%) male and 37 (49.3%) female in control group. Male to female ratio in case and control group was 1.08:1:1.02:1. The mean age of study population (Mean ± SD) in cases was 48.21± 13.76 and in controls 45.93± 12.03 years. (p=0.2818). In the study of Klara Paudel [8] had mean age of case was 47.2 ± 15.6 years & Joan C. Lo et al [9] had mean age was 48.7 ± 18.9 years. Duration of symptoms of CKD ranged from 3 months to 48 months. The mean duration was 10.23 ± 7.87 and Median being 8 months. Kayima et al [10] had mean duration of symptoms was 14.2 ± 9.9 months.

According to eGFR, three groups were made in our study; of which there were maximum 51 people (68%) having GFR <15 ml/min/1.73m². 18 (24%) with GFR between 15-29 ml/min/1.73m² and 6 (8%) people with eGFR ≥30 ml/min/1.73m²

Thyroid function tests in cases and controls

| Thyroid Function | Cases (Mean) | Controls (Mean) | P value |
|------------------------------|--------------|-----------------|---------|
| Total T ₃ (ng/dl) | 69.58 ± 38.2 | 113.9 ± 15.4 | <0.001 |
| Total T ₄ (g/dl) | 6.07 ± 2.55 | 7.54 ± 1.38 | <0.001 |
| TSH (IU/ml) | 7.42 ± 4.25 | 2.13 ± 0.87 | <0.001 |

Comparison of thyroid function between cases and controls

Overt Hypothyroidism found in 32 patients (42.66%) and Hyperthyroidism in 2 patients (2.67%) in CKD patients and controls were normal.

Correlation of T3, T4 and TSH with eGFR in cases and controls

With the increasing severity of CKD (decreased in eGFR), Overt hypothyroidism

also increase from 6.66% in stage-III, 38.8% in stage IV CKD to 47.06% in stage-V CKD and Hyperthyroidism present in stage-III and IV CKD patients. Chandra A [11] observed that the prevalence of overt hypothyroidism was increases from 6.28% in stage II, 11.87% in stage-III, and 40.68% in stage IV to 40.68% in stage V.

Thyroid dysfunction (n=75)

| S. No. | Type of Thyroid Dysfunction | No. of patients | % of thyroid dysfunction |
|--------|---|-----------------|--------------------------|
| 1 | Low T ₃ (Normal TSH) | 16 | 21.33 % |
| 2 | Hypothyroidism (Low T ₃ , Low T ₄ , high TSH) | 32 | 42.66 % |
| 3 | Sub clinical Hypothyroidism (Normal T ₃ ; High TSH) | 10 | 13.33 % |
| 4 | High T ₄ | 2 | 2.66% |
| 5 | High T ₃ | 2 | 2.66% |
| 6 | Hyperthyroidism (low TSH, High T ₃ &T ₄) | 2 | 2.66 % |

We found 60 out of 75 patients (80%), to be having some sort of thyroid dysfunction. Chandra A [11] and Prajapati et al [7] had 70% thyroid dysfunction in CKD patients. In our study, Low T₃ syndrome (Low T₃, Normal TSH) was found in 16 patients (21.33 %). Sub-clinical hypothyroidism (Raised TSH, Normal T₃,

T₄) was found in 10 patients (13.33 %). Overt Hypothyroidism (Low T₃ or low T₄, raised TSH) was found in 32 (42.66 %) patients.

Thus, in our study a positive, significant correlation is seen with the fall in eGFR and the level of T3 and TSH. [3,12]

Correlation of eGFR with thyroid function tests

| TFT | eGFR | | | P* _{1/2} | P* _{1/3} | P* _{2/3} | Anova p-value |
|-----|--------------|---------------|---------------|-------------------|-------------------|-------------------|---------------|
| | < 15 (51) | 15 – 29 (18) | ≥30 (6) | | | | |
| TT3 | 60.19± 25.60 | 78.19 ± 39.30 | 123.58± 72.21 | 0.181 | 0.020 | 0.001 | <0.001 |
| TT4 | 5.68± 2.27 | 6.42 ± 2.73 | 8.26 ± 3.43 | 0.851 | 0.362 | 0.057 | >0.05 |
| TSH | 8.38 ± 3.84 | 6.29 ± 4.37 | 2.59 ± 3.70 | 0.173 | 0.156 | 0.003 | <0.002 |

Symptoms and sign prevalence in other and overt hypothyroid groups in cases

| Symptoms | Other* (n = 41) | Overt Hypothyroid (n = 32) | P-value | OR |
|----------------------|-----------------|----------------------------|---------|------|
| Dry Skin | 11/41 (26.8%) | 23/32 (71.8%) | <0.05 | 0.14 |
| Fatigue | 24/41 (58.5%) | 20/32 (62.5%) | >0.05 | 0.85 |
| Cold intolerance | 5/41 (12.1%) | 19/32 (59.3%) | <0.001 | 0.10 |
| Tingling sensation | 7/41 (17.07%) | 17/32 (53.1%) | <0.05 | 0.18 |
| Hair loss | 8/41 (19.5%) | 7/32 (21.8%) | >0.05 | 0.87 |
| Constipation | 12/41 (29.26%) | 13/32 (40.6%) | >0.05 | 0.60 |
| Physical examination | Other* | Overt Hypothyroid | P-value | OR |
| Periorbital Edema | 31/41 (75.6%) | 31/32 (96.8%) | <0.05 | 0.10 |
| Pleural Effusion | 11/41 (26.82%) | 17/32 (53.1%) | <0.05 | 0.32 |
| Ascites | 14/41 (34.14%) | 23/32 (71.8%) | <0.01 | 0.20 |
| Pericardial Effusion | 6/41 (14.6%) | 15/32 (46.8%) | <0.01 | 0.19 |

Comparison of biochemical parameters in healthy controls and chronic kidney disease patients

| S. No. | Parameters | Case | Control | Sig. (2 tailed) P value |
|--------|-----------------------------------|--------------|---------------|-------------------------|
| 1 | Urea (mg/dl) | 183.7± 85.53 | 22.83 ± 5.06 | <0.001 |
| 2 | Creatinine (mg/dl) | 6.74 ±3.84 | 0.70±0.09 | <0.001 |
| 3 | Total protein (g/dl) | 6.15 ±0.68 | 7.02 ±0.60 | <0.001 |
| 4 | Albumin (g/dl) | 3.23 ±0.50 | 4.03 ±0.48 | <0.001 |
| 5 | Sodium (meq/l) | 133.2 ±4.72 | 140.4 ±4.07 | <0.001 |
| 6 | Potassium (meq/l) | 5.33 ±0.62 | 4.23 ±0.31 | <0.001 |
| 7 | Calcium (mg/dl) | 8.62 ±0.91 | 9.55 ±0.67 | <0.001 |
| 8 | eGFR (ml/min/1.73m ²) | 12.25 ±8.22 | 114.95 ±23.41 | <0.001 |

Comparison of lipid parameters between cases and controls

| S. No. | Parameters | Case | Control | Sig. (2 tailed) P value |
|--------|-------------------|----------------|----------------|-------------------------|
| 1 | Serum Cholesterol | 183.4 ± 48.1 | 136.07 ± 20.29 | <0.001 |
| 2 | HDL | 39.76 ± 8.53 | 41.67 ± 4.722 | 0.092 |
| 3 | LDL | 105.77 ± 36.28 | 99.63 ± 24.07 | 0.224 |
| 4 | TG | 142.51 ± 59.85 | 116.96 ± 30.95 | <0.001 |
| 5 | VLDL | 26.65 ± 9.83 | 25.13 ± 6.88 | 0.275 |
| 6 | S.CHL/HDL | 4.57 ± 0.94 | 3.72 ± 0.91 | <0.001 |

DISCUSSION

There was significant reduction of serum T3 level, T4 level and elevation of TSH level in cases (CKD patients) as compared to controls (normal healthy patients). [13,14,16,17] In comparison, TT₃ in 40 (53.33%) patients are more reduced than TT₄ 23 (30.67%) patients. Thyroid Stimulating Hormone levels were raised in 42 (56 %). Thus altered thyroid function was seen in 60 patients of which overt Hypothyroidism (Low T₃ or low T₄, raised TSH) was found in 32 (42.66%) patients, Low T₃ syndrome (Low T₃, Normal TSH) was found in 16 patients (21.33%) and Sub-clinical hypothyroidism (Raised TSH, Normal T₃, T₄) was found in 10 patients (13.33 %). A significant correlation was found between fall in eGFR and the elevated level of TSH (p<0.002) and decreased in serum TT₃ level (p<0.001) but not with TT₄ level.

Among the symptoms, Cold intolerance (p<0.001), Dry Skin, Tingling sensation were significantly (P<0.05) more prevalent in overt hypothyroid group as compared to other. [8,18]

On Clinical Examination, Periorbital edema (p<0.05), Pleural Effusion (p<0.05), Ascites and Pericardial Effusion were significantly (P<0.01) more prevalent in hypothyroid group as compared to other. [8,18]

Hypercholesterolemia and hypertriglyceridemia and S.CHL/HDL level were significantly higher (p value<0.01) in cases as compared to normal healthy control. [15]

Total protein, albumin, sodium, Calcium levels were significantly decreased in the CKD patients (P<0.001) and potassium level significantly increase in CKD patients (P<0.001).

CONCLUSIONS

The present study finds thyroid dysfunction to be very common in CKD patients and reveals the significant association between CKD progression and thyroid dysfunction and mean of T3, T4 decreases and TSH increases significantly in cases as compared to controls.

REFERENCES

1. Ramírez G, Jubiz W, Gutch CF, et al. Thyroid abnormalities in renal failure. A study of 53 patients on chronic hemodialysis. *Ann Intern Med* 1973; 79:500.
2. J. J. Carrero , A. R. Qureshi , J. Axelsson ,M.I. Yilmaz, et al. Clinical and biochemical implications of low thyroid hormone levels (total and free forms) in euthyroid patients with chronic kidney disease. *J Intern Med.* 2007 Dec;262(6): 690-701.
3. Hollowell JG, Staehling NW, Flanders WD, Hannon WH, Gunter EW, Spencer CA, Braverman LE. Serum TSH, T4, and thyroid antibodies in the United States population (1988 to 1994): National Health and Nutrition Examination Survey (NHANES III). *The Journal of Clinical Endocrinology & Metabolism.* 2002 Feb 1;87(2):4
4. Kher V. ESRD in developing countries *Nephrology forum. Kidney Int* 2002;62(1):350-62.
5. Prabahar MR, Chandrasekaran V, Soundararajan P. Epidemic of chronic kidney disease in India-what can be done? *Saudi Journal of Kidney Diseases and Transplantation.* 2008 Sep 1;19(5): 847.
6. Reddy KS, Shah B, Varghese C, Ramadoss A. Responding to the threat of chronic diseases in India. *The Lancet.* 2005 Nov 18;366(9498):1744-9.
7. Prajapati P, Singh AP, Bendwal S. Correlation between severity of chronic

- kidney disease and thyroid dysfunction. Journal of the Indian Medical Association. 2013 Aug;111(8):514-6.
8. Paudel K. Prevalence and clinical characteristics of hypothyroidism in a population undergoing maintenance hemodialysis. Journal of clinical and diagnostic research: JCDR. 2014 Apr;8(4):MC01.
 9. Lo JC, Chertow GM, Go AS, Hsu CY. Increased prevalence of subclinical and clinical hypothyroidism in persons with chronic kidney disease. Kidney international. 2005 Mar 31;67(3):1047-52.
 10. Kayima JK, Otieno LS, Gitau W, Mwai S. Thyroid hormone profiles in patients with chronic renal failure on conservative management and regular haemodialysis. East African medical journal. 1992 Jun;69(6):333-6.
 11. Chandra A. Prevalence of hypothyroidism in patients with chronic kidney disease: a cross-sectional study from North India. Kidney Research and Clinical Practice. 2016 Sep 30;35(3):165-8.
 12. Yoon, Yi Sun Jang, Jong Min Lee, HyeSoo Kim, Chul Woo Yang, Suk Young Kim, Hyeon Seok Hwang Kim EO. Unresolved Subclinical Hypothyroidism is Independently Associated with Progression of Chronic Kidney Disease. International Journal of Medical Sciences 2014;11(1):52-59.
 13. Shamsuddin M, Asmabi M. Chronic renal failure, Hypothyroidism, Hyperthyroidism, Thyroid Stimulating hormone (TSH). Comparative study of thyroid abnormalities with severity of chronic renal failure. 2015 May 10(9167).
 14. Rajeev G, Rayappa WD, Vijayalakshmi R, Swathi M, Kumar S. Evaluation of thyroid hormone levels in chronic kidney disease patients. Saudi Journal of Kidney Diseases and Transplantation. 2015 Jan 1;26(1):90.
 15. Khatiwada S, Rajendra KC, Gautam S, Lamsal M, Baral N. Thyroid dysfunction and dyslipidemia in chronic kidney disease patients. BMC endocrine disorders. 2015 Oct 29;15(1):1.
 16. Kumar R, Singh MP, Mehdi MD. Hospital based study on thyroid function in patients with chronic kidney disease in kosi region of bihar. International Journal of Recent Scientific Research IJRSR 2015 April Vol. 6, Issue, 4, pp.3412-3415.
 17. Lim VS, Fang VS, Katz AI, Refetoff S. Thyroid dysfunction in chronic renal failure: a study of the pituitary-thyroid axis and peripheral turnover kinetics of thyroxine and triiodothyronine. Journal of Clinical Investigation. 1977 Sep;60(3):522.
 18. Canaris GJ, Manowitz NR, Mayor G, Ridgway EC. The Colorado thyroid disease prevalence study. Archives of internal medicine. 2000 Feb 28;160(4): 526-34.

How to cite this article: Punekar J, Singh AA, Malav MK. Study of thyroid function in patients with chronic kidney disease. Int J Health Sci Res. 2017; 7(2):68-72.
