

Original Research Article

# The Role of Demographic Factors in Cognitive Impairments of Children with Autism Spectrum Disorder

S. Gopal Jee

Associate Professor & Head, Department of Psychology, DAV PG College (B. H.U.), Varanasi

## ABSTRACT

Autism Spectrum Disorder (ASD) is a clinically heterogeneous condition with a wide range of etiological factors and causing significant public health burden. The present study was designed to assess the cognitive ability and severity of autism spectrum disorder in context of various demographic factors. Twenty five children with autism spectrum disorder (18 mild and 7 moderate) and twenty five normal controls, , aged 4-11 years were selected from Indian Institute of cerebral palsy center, Varanasi. Indian Scale for Assessment of Autism (ISAA) and Child Autism Spectrum Quotient were used for assessment of cognitive ability. Results indicate that there is significant difference among mild, moderate and normal controls on Attention switching, attention detail, imagination and communication skills ( $F=101.41, 21.36, 161.97$  and  $493.97, p<0.01$  respectively). The current results also show sex, habitat and SES related differences in the cognitive profile of ASD children. There is significant correlation with demographic variable and cognitive measures in children with Autism spectrum disorder. The study provides evidence that demographic factors influence the cognitive functions in children with ASD.

**Key words:** Autism, cognition, gender, Region

## INTRODUCTION

Autism spectrum disorder (ASD) is a developmental disorder characterised by impaired social interaction and communication, and restricted and repetitive patterns of behaviour and interests (RRBIs) (American Psychiatric Association, 2013). The term “spectrum” refers to the wide range of symptoms, skills and level of impairments that children with ASD can have. These behavioural symptoms are thought to reflect underlying cognitive deficits/differences, which have been extensively researched (Brunsdon & Happe, 2014). It is well established that there is tremendous variability in outcome in autism. Long-term outcome studies have shown that while a majority of individuals exhibit poor to very poor outcomes, many individuals with autism go on to achieve adequate levels of academic, social, and

occupational functioning (Nordin & Gillberg, 1998; Sigman & Norman, 1999). In a previous study that followed children with autism from age 2 to 9, as many as 40% were found to have good outcomes based on language and cognitive scores (Stone, Turner, Pozdol, & Smoski, 2003).

ASD occur in approximately 1 to 2% of the population and are more frequently diagnosed in males, with a sex ratio of around 4:1 (Baird et al., 2006; Centers for Disease Control and Prevention, 2014). Considering individuals without an intellectual disability only, the sex ratio increases to 8-9:1 (Mandy et al., 2011; Scott et al., 2002), indicating that sex differences in the vulnerability for autism change as a function of IQ. However, others suggest this discrepancy in autism diagnosis may be overestimated and report overall sex ratios between 2.0 and 2.6:1 (Kim et al., 2011;

Mattila et al., 2011). This differential diagnosis has led to an expanding body of research evaluating sex differences in autism. Understanding sex differences in more detail is important for unpacking the complex aetiology of autism (Rutter et al., 2003). An evaluation of the research to date reveals a number of inconsistencies in the clinical autism phenotype across males and females. Some studies have indicated that autistic females display more severe social and communication difficulties compared to autistic males (Hartley and Sikora, 2009), while others have indicated that these difficulties are less severe in autistic women (McLennan et al., 1993), or report no sex differences (Wilson et al., 2016). Some studies suggest that autistic males show more stereotyped and repetitive behaviours than females (Hattier et al., 2011; Van Wijngaarden-Cremers et al., 2014) and that females on the spectrum display more socially acceptable special interests (Gould and Ashton-Smith, 2011). However, others argue that these interests and behaviours are similar across sex (Harrop et al., 2015). It has also been suggested that in order to receive a diagnosis, women are required to display more impairment in functioning than men (Dworzynski et al., 2012), yet other studies fail to confirm these findings when controlling for IQ (Holtmann et al., 2007; Pilowsky et al., 1998).

Geographic variation in the occurrence of ASD has been studied also, and geographical spatial clustering of ASD has recently been observed (Mazumdar et al. 2010, Van Meter et al. 2010). An association between ASD and urbanicity (i.e. higher risk of autism in urban versus rural districts) has been documented (Lauritsen et al. 2005; Williams et al. 2006; Rosenberg et al. 2009; Lai et al. 2012). Urbanicity may act as a proxy for other factors, e.g. better access to diagnostic and medical services or higher exposure to as yet unknown environmental risk factors for ASD in urban vs. rural areas.

Given the inconsistencies in previous research, it is important to study

sex and geographical variation in the children with autism. The evidences indicated that children with autism spectrum disorders in India have problems in various spheres of life like social relationships, language & communication problems and cognitive problems. However these areas of research have got least attention from researchers. The diversity of culture and habitat in India may bear on the manifestation of the autism phenotype. Geographical factors have been shown to influence autism prevalence in other countries. Particularly in the case of social communicative deficits toward the autism spectrum, symptoms manifest only against the background of cultural norms and therefore the very existence of some symptoms may be partially a function of culture. In view of the above research gap and controversies, the present study attempts to compare the impact of gender and habitat on social skill, attention switching, attention detail, communication and Imagination among normal, mild and moderate children with Autism spectrum.

### **Objectives**

1. To assess and compare the social skill, attention switching, attention detail, communication and Imagination among normal, mild and moderate children with Autism spectrum.
2. To compare the impact of Gender on the social skill, attention switching, attention detail, Communication and imagination among normal, mild and moderate children with Autism spectrum.
3. To assess the impact of Habitat on social skill, attention switching, attention detail, Communication and imagination among normal, mild and moderate children with Autism spectrum.
4. To explore the relationship between demographic factor (sex & habitat) and different Dimension of AQ.

## **MATERIALS AND METHODS**

### **Sample**

The sample comprised of 25 diagnosed children with autism spectrum disorder (18 mild and 7 moderate) screened on the basis of Indian Scale for Assessment of Autism (ISAA) and 25 matched controls (Age range 4-11 years) and performed Child Autism Spectrum Quotient. The Autistic children were taken from “Indian Institute of Cerebral Palsy Center Sigara, Varanasi. Children who were suffering from other category of Mental retardation and organic lesion were not included in this study.

**Procedure**

The participants were approached after taking permission from the respective center and consent from the parents of children. The parents of participant were given full information about the questionnaire and the process of research. The questions of the parents were also handling. In all process confidentiality will be maintained and parents of participant were briefed that the information they will be used only for research purpose.

**Tools**

1. **Indian Scale for Assessment of Autism (ISAA):** The ISAA is a 40 item scale divided into six domains- Social Relationship and Reciprocity (9 questions); Emotional Responsiveness (5 questions); Speech — Language and Communication (9 questions); Behavior Patterns (7 questions); Sensory Aspects (6 questions) and Cognitive Component (4 questions). The scores for the each item of ISAA range from 1-5 and total score range was 40-200. The lowest

score represents no symptoms or symptoms which were present only rarely, and the maximum score indicates the most severe presentation of AD.

2. **Hindi version of the Child Autism Spectrum Quotient:** This is a 50-item parent-report questionnaire developed to detect autistic traits in children 4–11 years of age (Auyeung et al). Higher scores indicate a greater number of autistic traits. The AQ-Child has shown good test–retest reliability, high sensitivity (95%) and high specificity (95%). The AQ-Child was originally designed with five subscales to assess various domains of functioning: social skills; attention to detail; attention switching; communication; and imagination. Each domain is assessed by 10 questions, giving a maximum attainable score of 150.

**RESULTS**

This study was carried out with the purpose of investigating the role of demographic factors in cognitive impairments of children with Autism spectrum disorder. The collected data was tabulated and was analyzed in the light of objectives framed. To compare the cognitive patterns among normal, mild and moderate children with Autism spectrum, one way ANOVA was applied to test the significance of severity difference in mean scores on these variables, which is displayed in table 1 & table 2, Low score on different domains of AQ is denoting more severity.

Table-1: Mean, SD and One way ANOVA among Normal, mild and moderately autistic children on different dimension of AQ.

Domains AQ	Autism stages	N	Mean	SD	F ratio
Social skill	Mild	18	20.67	5.099	170.668**
	Moderate	7	23.00	2.309	
	Severe	25	3.32	1.909	
Attention switching	Mild	18	17.33	3.614	101.415**
	Moderate	7	21.14	3.532	
	Severe	25	7.12	1.9223	
Attention detail	Mild	18	8.83	4.328	21.362**
	Moderate	7	7.71	3.904	
	Severe	25	16.24	4.166	
communication	Mild	18	23.72	4.012	493.976*
	Moderate	7	26.14	1.215	
	Severe	25	1.52	1.122	
imagination	Mild	18	20.67	4.058	161.975**
	Moderate	7	25.57	2.699	
	Severe	25	6.36	2.413	

\*\* P<.01

An inspection of table-1 indicates that the mean values of the normal children are lower than mildly and moderately autistic children on dimensions of Child Autism Spectrum Quotient. Higher scores indicate a greater number of autistic traits. Results of one way ANOVA, which compared the three groups depending on severity of autism, indicated that there were statistically significant differences between

the groups for Social skills ( $F_{2,47} = 170.67$ ,  $P < 0.01$ ), Attention switching ( $F_{2,47} = 101.41$ ,  $P < 0.01$ ), Attention detail ( $F_{2,47} = 21.36$ ,  $P < 0.01$ ), communication ( $F_{2,47} = 493.97$ ,  $P < 0.01$ ) and imagination ( $F_{2,47} = 161.97$ ,  $P < 0.01$ ). In order to find out the differences in different domains of Autism Quotient by male and female, t test was computed, which is displayed in Table-2

**Table- 2: Mean, SD and t values for Autism Quotient across gender**

Autism Quotient	Gender	N	Mean	Std. deviation	t value
Social skill	Boys	33	14.82	9.879	2.685**
	Girls	17	7.47	7.543	
Attention switching	Boys	33	14.45	6.638	2.749**
	Girls	17	9.47	4.745	
Attention detail	Boys	33	11.18	5.736	2.156*
	Girls	17	14.71	4.909	
communication	Boys	33	15.88	11.518	2.559**
	Girls	17	7.29	10.646	
imagination	Boys	33	16.24	8.913	2.444*
	Girls	17	10.24	6.666	
Total ASQ	Boys	33	72.58	31.453	2.633**
	Girls	17	49.18	26.085	

\* p< .05; \*\* p< .01

Table-2 represents the result for autism quotient between boys and girls groups. Boys scored high on all domains of AQ than girls (accept attention details) which indicates a greater number of autistic traits in boys. The t values for gender

difference on different dimensions of AQ are significant at .01 level, which indicates that boys and girls differ in their social skills, attention switching, attention detail, communication and imagination skills.

**Table-3: Mean, SD and t values for Autism Quotient across habitat**

Autism Quotient	Habitat	N	Mean	Std. deviation	t value
Social skill	urban	29	6.03	7.144	8.27**
	rural	21	21.00	4.909	
Attention switching	urban	29	8.83	4.759	7.22**
	rural	21	18.19	4.167	
Attention detail	urban	29	14.83	5.278	4.130**
	rural	21	9.00	4.382	
communication	urban	29	5.03	9.018	9.038**
	rural	21	23.90	3.687	
imagination	urban	29	8.90	6.878	7.371**
	rural	21	21.52	4.423	
Total ASQ	urban	29	43.06	23.579	8.951**
	rural	21	93.62	11.561	

\*\* p< .01

Table-3 shows the result for autism quotient between urban and rural groups. Children resided in rural areas scored high on all domains of AQ than resided in urban areas (accept attention details) which

indicates a greater number of autistic traits in rural area. The t values for geographical differences on different dimensions of AQ are significant at .01 level, which indicates the higher risk of autism in urbanicity.

**Table-4: Correlation coefficient between demographic variables and all dimensions of AQ**

Demo variable	Social skill	Attention switching	Attention detail	Communication	Imagination
Gender	-.361**	-.369**	.297*	-.347*	-.355*
Habitat	.767**	.722**	-.512**	.794**	.791**

\*p<0.05, \*\*P<0.01

Table -4 reveals the relationship between Gender, Location and domains of AQ. It is evident from the table 3 that the coefficient of correlation between Social skills, attention switching, communication and imagination with gender is negative and significant at .05 level and the coefficient of correlation between Domains of AQ with their habitat is positive and also significant at .01 level.

## DISCUSSION

The present study was aimed to find out the differences between children with mild ASD, moderate ASD and normal controls on AQ. This study also aimed to find out any differences with regards to gender and region. The major findings of this study are that mildly and moderately autistic children are unable to communicate and lack of social skills. Children with ASD may have difficulty developing language skills and understanding what others say to them. They also often have difficulty communicating nonverbally, such as through hand gestures, eye contact, and facial expressions. ASD affects people in different ways and can range from mild to severe so moderately autistic children more effected than mildly and normal controls. Autism spectrum disorders (ASD) is characterized by deficits in two core domains: deficits in social communication and social interaction; restricted repetitive patterns of behavior, interests and activities (APA,2013).

It is well established that females are much less likely to be diagnosed with an autism disorder (ASD) than males, with an approximate ratio of 5:1 (e.g., Centers for Disease Control and Prevention, 2012) despite manifesting equivalent levels of autistic traits as assessed by the Childhood Autism Spectrum Test (Dworzynski, Ronald, Bolton & Happé, 2012). Results of the present study reveals that children resided in rural areas are more effected which support the other study (Mazumdar et al. 2010). An association between ASD and urbanicity (i.e. higher risk of autism in

urban versus rural districts) has been documented (Lauritsen et al. 2005; Williams et al. 2006; Rosenberg et al. 2009; Lai et al. 2012). Urbanicity may act as a proxy for other factors, e.g. better access to diagnostic and medical services or higher exposure to as yet unknown environmental risk factors for ASD in urban vs. rural areas. Autism spectrum disorder (ASD) diagnoses are made based on a pattern of behavioral symptoms, yet a growing body of research indicates that when, and indeed whether, an individual receives a diagnosis of ASD is influenced by demographic factors including sex, habitat, ethnicity, socioeconomic status (SES), and parental education level.

## REFERENCES

- American Psychiatric Association (2013). *Diagnostic and Statistical Manual of Mental Disorders 5th edition (DSM-5)*. Washington, DC: American Psychiatric Association
- Baird G, Simonoff E, Pickles A, et al. (2006) Prevalence of disorders of the autism spectrum in a population cohort of children in South Thames: the special needs and autism project (SNAP). *The Lancet* 368: 210–215.
- Brunson, V.E.A., & Happe, F. (2014). Exploring the ‘fractionation’ of autism at the cognitive level. *Autism: the International journal of research and practice*, 18,17-30
- Centers for Disease Control and Prevention (2014) Prevalence of autism spectrum disorder among children aged 8 years: autism and developmental disabilities monitoring network, 11 sites, United States, 2010. *MMWR Surveillance Summaries* 63: 1–21.
- Dworzynski K, Ronald A, Bolton P, et al. (2012) How different are girls and boys above and below the diagnostic threshold for autism spectrum disorders? *Journal of the American Academy of Child and Adolescent Psychiatry* 51: 788–797

- Gould J and Ashton-Smith J (2011) Missed diagnosis or misdiagnosis? Girls and women on the autism spectrum. *Good Autism Practice* 12: 34–41.
- Hattier MA, Matson JL, Tureck K, et al. (2011) The effects of gender and age on repetitive and/or restricted behaviors and interests in adults with autism spectrum disorders and intellectual disability. *Research in Developmental Disabilities* 32: 2346–2351.
- Harrop C, Gulsrud A and Kasari C (2015) Does gender moderate core deficits in ASD? An investigation into restricted and repetitive behaviors in girls and boys with ASD. *Journal of Autism and Developmental Disorders* 45: 3644–3655.
- Holtmann M, Bölte S and Poustka F (2007) Autism spectrum disorders: sex differences in autistic behaviour domains and coexisting psychopathology. *Developmental Medicine & Child Neurology* 49: 361–366.
- Larsson HJ, Eaton WW, Madsen KM, Vestergaard M, Olesen AV, Agerbo E, Schendel D, Thorsen P and Mortensen PB (2005) Risk factors for autism: perinatal factors, parental psychiatric history, and socioeconomic status. *American Journal of Epidemiology* 161(10): 916–925.
- Kim YS, Leventhal BL, Koh YJ, et al. (2011) Prevalence of autism spectrum disorders in a total population sample. *American Journal of Psychiatry* 168: 904–912.
- Lauritsen MB.(2013). Autism spectrum disorders. *European Child and Adolescent Psychiatry*. 22
- Mazumdar S, King M, Liu KY, Zerubavel N, Bearman P.(2010). The spatial structure of autism in California, 1993–2001. *Health Place*. 16:539–546.
- Mandy W, Charman T, Gilmour J, et al. (2011) Toward specifying pervasive developmental disorder-not otherwise specified. *Autism Research* 4: 121–131.
- Mattila ML, Kielinen M, Linna SL, et al. (2011) Autism spectrum disorders according to DSM- IV-TR and comparison with DSM-5 draft criteria: an epidemiological study. *Journal of the American Academy of Child and Adolescent Psychiatry* 50: 583–592.
- Rai D, Lewis G, Lundberg M, Araya R, Svensson A, Dalman C, Carpenter P and Magnusson C (2012) Parental socioeconomic status and risk of offspring autism spectrum disorders in a Swedish population-based study. *Journal of the American Academy of Child and Adolescent Psychiatry* 51(5): 467–476.
- Rosenberg RE, Daniels AM, Law JK, Law PA, Kaufmann WE. (2009). Trends in autism spectrum disorder diagnoses: 1994–2007. *Journal of Autism and Developmental Disorders*. 39:1099–1111.
- Rutter M, Caspi A and Moffitt TE (2003) Using sex differences in psychopathology to study causal mechanisms: unifying issues and research strategies. *Journal of Child Psychology and Psychiatry* 44: 1092–1115.
- Scott FJ, Baron-Cohen S, Bolton P, et al. (2002) Brief report: prevalence of autism spectrum conditions in children aged 5–11 years in Cambridgeshire, UK. *Autism* 6: 231–237.
- Sigman, M., & Norman, K. (1999). Continuity and change in the development of children with autism. In S. Broman & J. Fletcher (Eds.), *The changing nervous system: Neurobehavioral consequences of early brain disorders* (pp. 274–291). New York: Oxford University Press.
- Stone, W. L., Turner, L. M., Pozdol, S. L., & Smoski, M. J. (2003, April). Changes in diagnostic and developmental features from age 2 to age 9 in children with ASD. Paper presented at the Biennial Meeting of the Society for Research in Child Development, Tampa, FL.

- Van Meter KC, Christiansen LE, Delwiche LD, Azari R, Carpenter TE, Hertz-Picciotto I.(2010) Geographic distribution of autism in California: A retrospective birth cohort analysis. *Autism Research*. 3;19-29.
- Van Wijngaarden-Cremers PJM, Van Eeten E, Groen WB, et al. (2014) Gender and age differences in the core triad of impairments in autism spectrum disorders: A systematic review and meta-analysis. *Journal of Autism and Developmental Disorders* 44: 627–635.
- Williams JG, Higgins JP, Brayne CE. (2006).Systematic review of prevalence studies of autism spectrum disorders. *Archives of Disease in Childhood*. 91:8–15.
- Wilson CE, Murphy CM, McAlonan G, et al. (2016) Does sex influence the diagnostic evaluation of autism spectrum disorder in adults? *Autism*. DOI: 10.1177/1362361315611381

How to cite this article: S. Gopal Jee. The role of demographic factors in cognitive impairments of children with autism spectrum disorder. *Int J Health Sci Res*. 2018; 8(10):26-32.

\*\*\*\*\*